

1. Funamoto M, Sunagawa Y, Katanasaka Y, Kato T, **Funada J**, Ajiro Y, Komiyama M, Akao M, Yasoda A, Yamakage H, Satoh-Asahara N, Wada H, Ikeda Y, Morimoto T, Hasegawa K. Effects of high-absorption curcumin for the prevention of hypertensive heart disease: a double-blind, placebo-controlled, randomized clinical study. *Eur Heart J (Open)*. 2022; 2: 1-9.
2. Yoshida T, Nakamura A, **Funada J**, Amino M, Shimizu W, Fukuzawa M, Watanabe S, Hayashi T, Yamashita T, Okumura K, Akao M. Efficacy and Safety of Edoxaban 15 mg According to Renal Function in Very Elderly Patients with Atrial Fibrillation: A Subanalysis of the ELDERCARE-AF Trial. *Circulation*. 2022 Mar 1; 145(9): 718–720.
3. Wada H, Shinozaki T, Suzuki M, Sakagami S, Ajiro Y, **Funada J**, Matsuda M, Shimizu M, Takenaka T, Morita Y, Yonezawa K, Matsubara H, Ono Y, Nakamura T, Fujimoto K, Ninomiya A, Kato T, Unoki T, Takagi D, Wada K, Wada M, Iguchi M, Yamakage H, Kusakabe T, Yasoda A, Shimatsu A, Kotani K, Satoh-Asahara N, Abe M, Akao M, Hasegawa K, for the EXCEED-J Study Investigators. Impact of Chronic Kidney Disease on the Associations of Cardiovascular Biomarkers with Adverse Outcomes in Patients with Suspected or Known Coronary Artery Disease: The EXCEED-J Study. *J Am Heart Assoc*. 2022; 11: e023464.
4. Iguchi M, Wada H, Shinozaki T, Suzuki M, Ajiro Y, Matsuda M, Koike A, Koizumi T, Shimizu M, Ono Y, Takenaka T, Sakagami S, Morita Y, Fujimoto K, Yonezawa K, Yoshida K, Ninomiya A, Nakamura T, **Funada J**, Kajikawa Y, Oishi Y, Kato T, Kotani K, Abe M, Akao M, Hasegawa K, for the PREHOSP-CHF Study Investigators. Soluble vascular endothelial growth factor receptor 2 and prognosis in patients with chronic heart failure. *ESC Heart Fail*. 2021 Oct; 8(5): 4187–4198.
5. Wada H, Suzuki M, Matsuda M, Ajiro Y, Shinozaki T, Sakagami S, Yonezawa K, Shimizu M, **Funada J**, Takenaka T, Morita Y, Nakamura T, Fujimoto K, Matsubara H, Kato T, Unoki T, Takagi D, Wada K, Wada M, Iguchi M, Masunaga N, Ishii M, Yamakage H, Kusakabe T,

- Yasoda A, Shimatsu A, Kotani K, Satoh-Asahara N, Abe M, Akao M, Hasegawa K; ANOX Study Investigators. Impact of smoking status on growth differentiation factor 15 and mortality in patients with suspected or known coronary artery disease: The ANOX Study. *J Am Heart Assoc.* 2020; Nov 10: e018217.
6. Wada H, Suzuki M, Matsuda M, Ajiro Y, Shinozaki T, Sakagami S, Yonezawa K, Shimizu M, **Funada J**, Takenaka T, Morita Y, Nakamura N, Fujimoto K, Matsubara H, Kato T, Unoki T, Takagi D, Wada K, Wada M, Iguchi M, Masunaga N, Ishii M, Yamakage H, Kusakabe T, Yasoda A, Shimatsu A, Kotani K, Satoh-Asahara N, Abe M, Akao M, Hasegawa K. Distinct Characteristics of VEGF-D and VEGF-C to predict mortality in patients with suspected or known coronary artery disease. *J Am Heart Assoc.* 2020; 9: e015761.
 7. Takakado M, Takata Y, Yamagata F, Yaguchi M, Hiasa G, Sato S, **Funada J**, Kawazu S, Osawa H. Simple and non-invasive screening method for diabetes based on myoinositol levels in urine samples collected at home. *BMJ Open Diabetes Res Care.* 2020; 8(1): e000984.
 8. Ono K, Wada H, Satoh-Asahara N, Inoue H, Uehara K, **Funada J**, Ogo A, Horie T, Fujita M, Shimatsu A, Hasegawa K; ABLE-MET Investigators. Effects of metformin on left ventricular size and function in hypertensive patients with type 2 diabetes mellitus: results of a randomized, controlled, multicenter, phase IV trial. *Am J Cardiovasc Drugs.* 2020; 20(3): 283–293.
 9. Wada H, Suzuki M, Matsuda M, Ajiro Y, Shinozaki T, Sakagami S, Yonezawa K, Shimizu M, **Funada J**, Takenaka T, Morita Y, Nakamura T, Fujimoto K, Matsubara H, Kato T, Unoki T, Takagi D, Ura S, Wada K, Iguchi M, Masunaga N, Ishii M, Yamakage H, Shimatsu A, Kotani K, Satoh-Asahara N, Abe M, Akao M, Hasegawa K; for the ANOX Study Investigators. VEGF-C and mortality in patients with suspected or known coronary artery disease. *J Am Heart Assoc.* 2018; 7: e010355.
 10. Fujii A, Inoue K, Nagai T, Nishimura K, Uetani T, Suzuki J, **Funada J**, Okura T, Higaki J,

- Ogimoto A. Clinical utility of atrial electromechanical conduction time measured with speckle tracking echocardiography after catheter ablation in patients with atrial fibrillation: A validation study with electroanatomical mapping. *Echocardiography*. 2016; 33(9): 1317-1325.
11. Fujii A, Inoue K, Nagai T, Uetani T, Nishimura K, Suzuki J, **Funada J**, Okura T, Higaki J, Ogimoto A. Clinical significance of peripheral endothelial function for left atrial blood stagnation in nonvalvular atrial fibrillation patients with low-to-intermediate stroke risk. *Circ J*. 2016; 80: 2117-2113.
 12. Yukumi S, Ichiki H, **Funada J**, Suzuki H, Morimoto M, Fujita T, Izumi N, Abe M. Postcardiac injury syndrome following vascular interventional radiofrequency ablation for paroxysmal atrial fibrillation. *Respiratory medicine case report*. 2015; 15: 89-91.
 13. Kurata A, Kawaguchi N, Kido T, Inoue K, Suzuki J, Ogimoto A, **Funada J**, Higaki J, Miyagawa M, Vembar M, Mochizuki T. Qualitative and quantitative assessment of adenosine triphosphate stress whole-heart dynamic myocardial perfusion imaging using 256-slice computed tomography. *PLoS One*. 2013 Dec 23; 8(12): e83950.
 14. Inoue K, Okayama H, Nishimura K, Nagai T, Suzuki J, Ogimoto A, Saito M, Yoshii T, Hiasa G, Sumimoto T, **Funada J**, Shigematsu Y, Hamada M, Higaki J. Impact of septal curvature on regional strain in patients with hypertrophic cardiomyopathy. *Circ J*. 2013; 77(4): 1040-1045.
 15. S Inaba, H Okayama, **J Funada**, H Higashi, M Saito, T Yoshii, G Hiasa, T Sumimoto, Y Takata, K Nishimura, K Inoue, A Ogimoto, J Higaki. Impact of type 2 diabetes on serial changes in tissue characteristics of coronary plaques -an integrated backscatter intravascular ultrasound analysis. *Eur Heart J. Cardiovasc Imaging*. 2012 Sep; 13(9):717-723.
 16. K Nishimura, H Okayama, K Inoue, M Saito, T Yoshii, G Hiasa, T Sumimoto, S Inaba, A Ogimoto, **J Funada**, J Higaki. Direct measurement of radial strain in the inner-half layer of

the left ventricular wall in hypertensive patients. *J Cardiol*. 2012 Jan; 59(1): 64-71.

17. K Inoue, H Okayama, K Nishimura, M Saito, T Yoshii, G Hiasa, T Sumimoto, S Inaba, J Suzuki, A Ogimoto, **J Funada**, J Higaki. Right ventricular septal pacing preserves global left ventricular longitudinal function in comparison with apical pacing: analysis is of speckle tracking echocardiography. *Circ J*. 2011; 75 1609-1615.
18. M Saito, H Okayama, T Yoshii, G Hiasa, T Sumimoto, S Inaba, K Nishimura, K Inoue, A Ogimoto, Y Shigematsu, **J Funada**, M Hamada M, J Higaki. J. Myocardial fibrosis attenuates the effect of cibenzoline on left ventricular diastolic function in patients with hypertrophic cardiomyopathy. *Cardiovasc Pharmacol*. 2011 Feb; 57(2): 207-212.
19. T Nagai, H Okayama, K Nishimura, K Inoue, J Suzuki, A Ogimoto, T Ohtsuka, G Hiasa, T Sumimoto, **J Funada**, J Higaki. Initial Japanese experience and long-term follow-up with a new active fixation coronary sinus lead, the StarFix 4195. *J Cardiol cases*. 2010; 1: e176-179.
20. K Nishimura, H Okayama, K Inoue, M Saito, T Yoshii, G Hiasa, T Sumimoto, S Inaba, A Ogimoto, T Ohtsuka, **J Funada**, Y Shigematsu, J Higaki. Usefulness of the MOSAIC (measurement of stenosis by aliasing coronary flow) method using transthoracic color Doppler echocardiography in unstable angina patients. *Int J Cardiol*. 2011 Sep 1; 151: 170-174.
21. M Saito, H Okayama, T Yoshii, G Hiasa, T Sumimoto, S Inaba, K Nishimura, K Inoue, A Ogimoto, T Ohtsuka, **J Funada**, Y Shigematsu, J Higaki. The differences in left ventricular torsional behavior between patients with hypertrophic cardiomyopathy and hypertensive heart disease. *Int J Cardiol*. 2011 Aug 4; 150: 301-306.
22. S Inaba, H Okayama, **J Funada**, H Hashida, G Hiasa, T Sumimoto, Y Takata, K Nishimura, K Inoue, A Ogimoto, T Ohtsuka, J Higaki. Relationship between smaller calcifications and lipid-rich plaques on integrated backscatter-intravascular ultrasound. *Int J Cardiol*. 2010;

145(2):347-348.

23. K Inoue, H Okayama, K Nishimura, A Ogimoto, T Ohtsuka, M Saito, G Hiasa, T Yoshii, T Sumimoto, **J Funada**, J Higaki. J. Right ventricular pacing from the septum avoids the acute exacerbation in left ventricular dyssynchrony and torsional behavior seen with pacing from the apex. *Am Soc Echocardiogr* 2010; 23(2): 195-200.
24. H Hashida, **J Funada**, N Morioka, T Iwata. The lurking potential of tangential forces: a case of an arteriovenous shunt developed by percutaneous coronary intervention for the septal branch. *J Cardiol* 2010; 1: e1-e5.
25. A Fujii, **J Funada**, N Morioka, H Hashida, T Iwata, T Higaki. A case of pulmonary stenosis after a repair for tetralogy of Fallot treated with percutaneous pulmonary valvuloplasty using a triple-balloon technique. *J Cardiol* 2010; 55: 130-134.